

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number
WO 01/40246 A1

- (51) International Patent Classification⁷: **C07H 19/16**,
A61K 31/7076, A61P 9/00, 25/00
- (21) International Application Number: **PCT/US00/32721**
- (22) International Filing Date: 1 December 2000 (01.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/454,136 3 December 1999 (03.12.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/454,136 (CON)
Filed on 3 December 1999 (03.12.1999)
- (71) Applicant (for all designated States except US): **CV THERAPEUTICS, INC.** [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ZABLOCKI, Jeff, A.** [US/US]; 580 Sleeper Avenue, Mountain View, CA 94040 (US). **PALLE, Venkata, P.** [IN/US]; 200 E. Dana Street, #F122, Mountain View, CA 94041 (US). **VARKHEDKAR, Vaibhav** [IN/US]; 142 S. Bernado Avenue #2, Sunnyvale, CA 94086 (US). **BELARDINELLI, Luiz** [US/US]; 704 University Drive, Menlo Park, CA 94025 (US).
- (74) Agent: **HUGHES, A., Blair**; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Suite 3200, Chicago, IL 60606 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- With international search report.
 - Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/40246 A1

(54) Title: PARTIAL OR FULL A₁ AGONISTS - N⁶ HETEROCYCLIC 5'-THIO SUBSTITUTED ADENOSINE DERIVATIVES

(57) Abstract: N⁶ heterocyclic 5' modified adenosine derivatives that are adenosine A₁ receptor partial or full agonists, and as such, are useful for modifying cardiac activity, modifying adipocyte function, treating central nervous system disorders, and treating diabetic disorders and obesity in mammals, and especially in humans.

WO 01/40246

PCT/US00/32721

TITLE: PARTIAL OR FULL A₁ AGONISTS - N⁶ HETEROCYCLIC 5'-THIO SUBSTITUTED ADENOSINE DERIVATIVES

BACKGROUND OF THE INVENTION

5 (1) **Field of the Invention**

This invention includes stable and useful drugs and pro-drugs that are N⁶ heterocyclic 5'-thio modified adenosine derivatives. The compositions of this invention are selective, partial or full adenosine A₁ receptor agonists, and as such, are useful for modifying cardiac activity, modifying adipocyte function, treating central nervous system disorders, and treating
10 diabetic disorders and obesity in mammals, and especially in humans.

(2) **Description of the Art**

There are at least two subtypes of adenosine receptors in the heart: A₁ and A_{2A}. Each subtype affects different physiological functions. The A₁ adenosine receptor mediates two distinct physiological responses. Inhibition of the cardiostimulatory effects of catecholamine
15 are mediated via the inhibition of adenylate cyclase, whereas the direct effects to slow the heart rate (HR) and to prolong impulse propagation through the AV node are due in great part to activation of I_{KAdo}. (B. Lerman and L. Belardinelli Circulation, Vol. 83 (1991), P 1499-1509 and J. C. Shryock and L. Belardinelli The Am. J. Cardiology, Vol. 79 (1997) P 2-10). Both, the anti-β-adrenergic action and direct depressant effects on SA and AV nodal function are
20 mediated by the A₁ receptor; there is no role for the A_{2A} receptor in this response to adenosine. A_{2A} receptors mediate the coronary vasodilatation caused by adenosine. Stimulation of the A₁ adenosine receptor accordingly shortens the duration and decreases the amplitude of the action potential of AV nodal cells, and hence prolongs the refractory period of the AV nodal cell. The consequence of these effects is to limit the number of impulses conducted from the atria
25 to the ventricles. This forms the basis of the clinical utility of A₁ receptor agonists for the treatment of supraventricular tachycardias, including termination of nodal re-entrant tachycardias, and control of ventricular rate during atrial fibrillation and flutter.

A clinical utility of A₁ agonists therefore is in the treatment of acute and chronic disorders of heart rhythm, especially those diseases characterized by rapid heart rate where the
30 rate is driven by abnormalities in the sinoatrial, atria, and AV nodal tissues. Such disorders include but are not limited to atrial fibrillation, supraventricular tachycardia and atrial flutter.

WO 01/40246

PCT/US00/32721

Exposure to A₁ agonists causes a reduction in the heart rate and a regularization of the abnormal rhythm thereby improving cardiovascular function.

A₁ agonists, through their ability to inhibit the effects of catecholamines, decrease cellular cAMP, and thus, should have beneficial effects in the failing heart where increased
5 sympathetic tone increases cellular cAMP levels. The latter has been shown to be associated with increased likelihood of ventricular arrhythmias and sudden death. All of the above concepts are discussed in reviews regarding the effects of adenosine on cardiac electrophysiology (see B. Lerman and L. Belardinelli *Circulation*, Vol. 83 (1991), P 1499-1509 and J. C. Shryock and L. Belardinelli, *Am. J. Cardiology*, Vol. 79 (1997) P 2-10).

10 A controversial area in the field of A₁ adenosine agonism is that the benefit of preconditioning of the heart prior to ischemia may be due to binding of adenosine to the A₁ receptor. Evidence for this hypothesis comes from a rabbit ischemia model wherein 2-chloro-N6-cyclopentyladenosine (CCPA) and R-PIA were administered prior to ischemia providing protection with respect to infarct size (J. D. Thornton et al. *Circulation* Vol. 85 (1992) 659-
15 665).

A₁ agonists, as a result of their inhibitory action on cyclic AMP generation, have antilipolytic effects in adipocytes that leads to a decreased release of nonesterified fatty acids (NEFA) (E. A. van Schaick et al *J. Pharmacokinetics and Biopharmaceutics*, Vol. 25 (1997) p 673-694 and P. Strong *Clinical Science* Vol. 84 (1993) p. 663-669). Non-insulin-dependent
20 diabetes mellitus (NIDDM) is characterized by an insulin resistance that results in hyperglycemia. Factors contributing to the observed hyperglycemia are a lack of normal glucose uptake and activation of skeletal muscle glycogen synthase (GS). Elevated levels of NEFA have been shown to inhibit insulin-stimulated glucose uptake and glycogen synthesis (D. Thiebaud et al *Metab. Clin. Exp.* Vol. 31 (1982) p 1128-1136 and G. Boden et al *J. Clin. Invest.* Vol. 93 (1994) p 2438-2446). The hypothesis of a glucose fatty acid cycle was
25 proposed by P. J. Randle as early as 1963 (P. J. Randle et al *Lancet* (1963) p. 785-789). A tenet of this hypothesis would be that limiting the supply of fatty acids to the peripheral tissues should promote carbohydrate utilization (P. Strong et al *Clinical Science* Vol. 84 (1993) p. 663-669).

30 The benefit of an A₁ agonist in central nervous disorders has been reviewed and the

WO 01/40246

PCT/US00/32721

content are included herein by reference (L. J. S. Knutsen and T. F. Murray In *Purinergic Approaches in Experimental Therapeutics*, Eds. K. A. Jacobson and M. F. Jarvis (1997) Wiley-Liss, N. Y., P 423-470). Briefly, based on experimental models of epilepsy, a mixed A_{2A} : A_1 agonist, metrifudil, has been shown to be a potent anticonvulsant against seizures
5 induced by the inverse benzodiazepine agonist methyl 6,7-dimethoxy-4-ethyl-beta-carboline-3-carboxylate (DMCM, H. Klitgaard Eur. J. Pharmacol. (1993) Vol. 224 p. 221-228). In other studies using CGS 21680, an A_{2A} agonist, it was concluded that the anticonvulsant activity was attributed to activation of the A_1 receptor (G. Zhang et al. Eur. J. Pharmacol. Vol. 255 (1994) p. 239-243). Furthermore, A_1 adenosine selective agonists have been shown to have
10 anticonvulsant activity in the DMCM model (L. J. S. Knutsen In *Adenosine and Adenine Nucleotides: From Molecular Biology to Integrative Physiology*; eds. L. Belardinelli and A. Pelleg, Kluwer: Boston, 1995, pp 479-487). A second area where an A_1 adenosine agonist has a benefit is in animal models of forebrain ischemia as demonstrated by Knutsen et al (J. Med. Chem. Vol. 42 (1999) p. 3463-3477). The benefit in neuroprotection is believed to be in part
15 due to the inhibition of the release of excitatory amino acids (ibid).

There are a number of full A_1 agonists disclosed in the prior art. However, the agonists disclosed are generally in the forms that are not useful in the mammalian body. Because useful forms of A_1 agonists may not always be stable, soluble or they may have other properties that make their incorporation into therapeutic dosage forms difficult, it is often
20 necessary to identify compositions that are more easily incorporated into therapeutic dosage forms in order to provide the desired therapeutic effect. Also, these agonists fail as useful therapeutics due to side effects caused by the non-selective stimulation of the A_1 adenosine receptor in all biologically available tissues and the desensitization of the desired response preempting their use as chronic agents. Therefore, there remains a need for specific and
25 selective A_1 agonists, precursors and/or pro-drugs that are converted in the body into useful therapeutic compositions.

30

WO 01/40246


PCT/US00/32721

SUMMARY OF THE INVENTION

In one aspect, this invention includes heterocyclic 5'-thio modified adenosine derivative compositions that are useful partial or full adenosine A₁ receptor agonists.

In another aspect, this invention includes pharmaceutical compositions including one or more heterocyclic 5'-thio modified adenosine derivative compositions that are well tolerated with few side effects.

In still another embodiment, this invention includes heterocyclic 5'-thio modified adenosine derivatives having the formula:



In yet another embodiment, this invention includes methods for administering compositions of this invention to mammals, and especially to humans, to stimulate coronary activity, to modify adipocyte function, to treat central nervous system disorders, and to treat diabetic disorders.


In a further embodiment, this invention is pharmaceutical compositions of matter comprising at least one composition of this invention and one or more pharmaceutical excipients.

WO 01/40246

PCT/US00/32721

DESCRIPTION OF THE CURRENT EMBODIMENT

This invention includes a class of heterocyclic 5'-thio modified adenosine derivatives having the formula having the formula:



5 wherein $X^1 = S, S(O), S(O)_2$;

wherein R^1 is a monocyclic or polycyclic heterocyclic group containing from 3 to 15 carbon atoms wherein at least one carbon atom is substituted with an atom or molecule selected from the group consisting of N, O, P and S-(O)₀₋₂ and wherein R^1 does not contain an epoxide group, and wherein R_2 is selected from the group consisting of hydrogen, halo, CF₃, and cyano; wherein R_3 and R_4 are independently selected from the group consisting of hydrogen, and -(CO)-R' and -(CO)-R'' wherein R' and R'' are independently selected from the group consisting of C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with 1 to 3 substituents independently selected from the group of halo, NO₂, heterocyclyl, aryl, heteroaryl, CF₃, CN, OR²⁰, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, SO₂NR²⁰COR²², SO₂NR²⁰CO₂R²², SO₂NR²⁰CON(R²⁰)₂, N(R²⁰)₂, NR²⁰COR²², NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, NR²⁰C(NR²⁰)NHR²³, COR²⁰, CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²², OC(O)R²⁰, C(O)OCH₂OC(O)R²⁰, and OCON(R²⁰)₂ and each optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with halo, NO₂, alkyl, CF₃, amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, NR²⁰COR²², NR²⁰SO₂R²², COR²⁰, CO₂R²⁰, CON(R²⁰)₂, NR²⁰CON(R²⁰)₂,

10
15
20

WO 01/40246

PCT/US00/32721

OC(O)R^{20} , $\text{OC(O)N(R}^{20})_2$, SR^{20} , S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, CN , or OR^{20} ;

wherein R_5 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with 1 to 3 substituents independently selected from the group consisting of halo, alkyl, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{S(O)}_3\text{R}^{20}$, S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, $\text{SO}_2\text{NR}^{20}\text{COR}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CON(R}^{20})_2$, $\text{P(O)(OR}^{20})_2$, $\text{N(R}^{20})_2$, $\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON(R}^{20})_2$, $\text{NR}^{20}\text{C(NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON(R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{OCONR}^{20}\text{SO}_2\text{R}^{22}$, OC(O)R^{20} , $\text{C(O)OCH}_2\text{OC(O)R}^{20}$, and $\text{OCON(R}^{20})_2$ and wherein optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, $\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, COR^{20} , CO_2R^{20} , $\text{CON(R}^{20})_2$, $\text{NR}^{20}\text{CON(R}^{20})_2$, OC(O)R^{20} , $\text{OC(O)N(R}^{20})_2$, SR^{20} , S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, CN , or OR^{20} ;

wherein R^{20} is a member selected from the group consisting of H, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl are optionally substituted with 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN , O-C_{1-6} alkyl, CF_3 , aryl, and heteroaryl; and

R^{22} is a member selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl are optionally substituted with 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN , O-C_{1-6} alkyl, CF_3 , and heteroaryl.

In preferred compositions, $\text{X}^1=\text{S}$ or SO_2 ; R_2 is a hydrogen; R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl and, more preferably, R_3 and R_4 are each hydrogen; R_5 is selected from the group consisting of C_{1-8} alkyl, and aryl wherein alkyl, and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, $\text{NR}^{20}\text{CON(R}^{20})_2$, CO_2R^{20} , $\text{CON(R}^{20})_2$, and

WO 01/40246

PCT/US00/32721

wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN, and OR^{20} ; R_{20} is selected from the group consisting of H, C_{1-6} alkyl; and R_{22} is selected from the group consisting of C_{1-6} . In the above compositions, R_5 is more preferably selected from the group consisting of C_{1-8} alkyl, and aryl wherein alkyl, and
 5 aryl are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, CF_3 , and OR^{20} .

In more preferred compositions, $\text{X}^1=\text{S}$ or SO_2 ; R_2 is a hydrogen; R_3 and R_4 are independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl
 10 which alkyl are optionally substituted with 1 substituent selected from the group consisting of aryl, CF_3 , CN, OR^{20} , $\text{N}(\text{R}^{20})_2$, and wherein each optional aryl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 ; R_5 is C_{1-8} alkyl, wherein alkyl, is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, CO_2R^{20} ,
 15 $\text{CON}(\text{R}^{20})_2$, wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN, and OR^{20} ; R^{20} is selected from the group consisting of H, C_{1-6} alkyl; and R_{22} is selected from the group consisting of C_{1-6} . In the above compositions, R_5 is more preferably C_{1-8} alkyl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of aryl, heteroaryl, OR^{20} ,
 20 $\text{S}(\text{O})\text{R}^{22}$, CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN, and OR^{20} , and R_5 is even more preferably C_{1-8} alkyl that is optionally substituted with 1 substituent selected from the group consisting of CO_2R^{20} , and $\text{CON}(\text{R}^{20})_2$, and R_5 is even more preferably C_{1-6} alkyl and most preferably methyl or ethyl or isopropyl. Also in the above compositions, R_3 and R_4 are
 25 more preferably each hydrogen and R_{20} is more preferably selected from the group consisting of H, and methyl.

In another class of preferred compositions, R_2 is a hydrogen; R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein each R' and R'' are independently selected from the group consisting of C_{1-6} alkyl,
 30 and aryl, which alkyl and aryl are optionally substituted with from 1 to 2 substituents

WO 01/40246

PCT/US00/32721

independently selected from the group of halo, NO₂, aryl, CF₃, CN, OR²⁰, N(R²⁰)₂, S(O)R²², SO₂R²², and wherein each optional aryl substituent is further optionally substituted with halo, NO₂, alkyl, CF₃; R₅ is selected from the group consisting of, aryl, and heteroaryl, wherein aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected

5 from the group consisting of halo, alkyl, aryl, heteroaryl, CF₃, CN, OR²⁰, SR²⁰, N(R²⁰)₂, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, CO₂R²⁰, CON(R²⁰)₂, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF₃, CO₂R²⁰, CON(R²⁰)₂, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, CN, or OR²⁰; R²⁰ is selected from the group consisting of H, C₁₋₆ alkyl, and aryl, which alkyl and aryl are optionally substituted with

10 1 substituent selected from halo, alkyl, mono- or dialkylamino, CN, O-C₁₋₆ alkyl, CF₃; and R²² is selected from the group consisting of C₁₋₆ alkyl and aryl, which alkyl and aryl are optionally substituted with 1 substituent selected from halo, alkyl or CN, O-C₁₋₆ alkyl, and CF₃. In the above compositions, X¹ is preferably S; R₃ and R₄ are more preferably hydrogen; R₅ is more preferably selected from the group consisting of, aryl, and heteroaryl, wherein aryl, and

15 heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF₃, CN, OR²⁰, SR²⁰, CO₂R²⁰, CON(R²⁰)₂. Even more preferably R₅ is aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, CF₃, OR²⁰, CO₂R²⁰, CON(R²⁰)₂. And most preferably, R₅ is phenyl that is optionally substituted with a substituent selected from the

20 group consisting of methoxy, chloro, fluoro, methyl, and trifluoromethyl. In the compounds above, R²⁰ is preferably selected from the group consisting of H, C₁₋₃ alkyl and most preferably H or methyl while R₂₂ is preferably selected from the group consisting of C₁₋₆ alkyl.


In the compositions of this invention, R₁ is preferably mono or polysubstituted with one or more compounds selected from the group consisting of halogen, oxo, hydroxyl, lower

25 alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof. More preferably, R₁ is a monocyclic, bicyclic, or tricyclic cycloalkyl group containing from 3 to 15 carbon atoms wherein at least one carbon atom is substituted with an atom or molecule selected from the group consisting of O or S-(O)₀₋₂. Some examples

30 of preferred R₁ moieties include:

WO 01/40246

PCT/US00/32721



wherein R₁['], R₁^{''}, R₁^{'''}, and R₁^{''''} may each individually be selected from the group halogen, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, and cyano, and X is O, or S (-O)_{0.2}, alternately, R₁^{'''} and R₁^{''''} may be a single oxygen atom. More preferably, R₁['], R₁^{''}, R₁^{'''}, and R₁^{''''} are each individually selected from the group hydrogen, lower alkyl, and substituted lower alkyl. In the compositions above, each R is individually selected from the group consisting of H, lower alkyl, and substituted lower alkyl and wherein X is O, or S (-O)_{0.2}.

- Most preferred compounds of this invention include, 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-(methylthiomethyl)oxolane-3,4-diol ; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(Ethylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(Methylethylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-(phenylthiomethyl)oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(4-Methoxyphenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(4-chlorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(4-fluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(4-methylphenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(4-(trifluoromethyl)phenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(2-Methoxyphenylthio)methyl]oxolane-3,4-diol; (5-{6-[[[(3R)oxolan-3-yl]amino]purinyl-9-yl]}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)(ethylsulfonyl)methane; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(2,4-difluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[[[(3R)oxolan-3-yl]amino]purin-9-yl]}(4S,5S,2R,3R)-5-[(2,6-

WO 01/40246

PCT/US00/32721

dichlorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-[(3-fluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol; 5-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(2S,3R,4R,5R)-4-acetyloxy-2-
 5 [(fluorophenylthio)methyl]oxolan-3-yl acetate; Methyl 2[(5-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]benzoate; {2[(5-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]phenyl}-N-methylcarboxamidebenzoate; 2-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(benzoxazol-2-ylthiomethyl)oxolane-3,4-diol; 2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-[(1-methylimidazol-2-yl-thio)methyl]oxolane-3,4-diol; 2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(pyrimidine-2-ylthiomethyl)oxolane-3,4-diol; 2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(2-pyridylthiomethyl)oxolane-3,4-diol; 2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(4-pyridylthiomethyl)oxolane-3,4-diol; and 5-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(2S,3R,4R,5R)-4-acetyloxy-2-[(4-fluorophenylthio)methyl]oxolan-3-yl]acetate.
 15

The following definitions apply to terms as used herein.

“Halo” or “Halogen” - alone or in combination means all halogens, that is, chloro (Cl), fluoro (F), bromo (Br), iodo (I).

“Hydroxyl” refers to the group -OH.

20 “Thiol” or “mercapto” refers to the group -SH.

“Alkyl” - alone or in combination means an alkane-derived radical containing from 1 to 20, preferably 1 to 15, carbon atoms (unless specifically defined). It is a straight chain alkyl, branched alkyl or cycloalkyl. Preferably, straight or branched alkyl groups containing from 1-15, more preferably 1 to 8, even more preferably 1-6, yet more preferably 1-4 and
 25 most preferably 1-2, carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl and the like. The term “lower alkyl” is used herein to describe the straight chain alkyl groups described immediately above. Preferably, cycloalkyl groups are monocyclic, bicyclic or tricyclic ring systems of 3-8, more preferably 3-6, ring members per ring, such as cyclopropyl, cyclopentyl, cyclohexyl, adamantyl and the like. Alkyl also includes a straight chain or
 30 branched alkyl group that contains or is interrupted by a cycloalkyl portion. The straight

WO 01/40246

PCT/US00/32721

chain or branched alkyl group is attached at any available point to produce a stable compound.

Examples of this include, but are not limited to, 4-(isopropyl)-cyclohexylethyl or 2-methyl-cyclopropylpentyl. A substituted alkyl is a straight chain alkyl, branched alkyl, or cycloalkyl group defined previously, independently substituted with 1 to 3 groups or substituents of halo,

- 5 hydroxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, acyloxy, aryloxy, heteroaryloxy, amino optionally mono- or di-substituted with alkyl, aryl or heteroaryl groups, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocyclyl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups, alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, or the like.

- “Alkenyl” - alone or in combination means a straight, branched, or cyclic hydrocarbon containing 2-20, preferably 2-17, more preferably 2-10, even more preferably 2-8, most preferably 2-4, carbon atoms and at least one, preferably 1-3, more preferably 1-2, most preferably one, carbon to carbon double bond. In the case of a cycloalkyl group, conjugation
- 15 of more than one carbon to carbon double bond is not such as to confer aromaticity to the ring.

- Carbon to carbon double bonds may be either contained within a cycloalkyl portion, with the exception of cyclopropyl, or within a straight chain or branched portion. Examples of alkenyl groups include ethenyl, propenyl, isopropenyl, butenyl, cyclohexenyl, cyclohexenylalkyl and the like. A substituted alkenyl is the straight chain alkenyl, branched alkenyl or cycloalkenyl
- 20 group defined previously, independently substituted with 1 to 3 groups or substituents of halo, hydroxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, acyloxy, aryloxy, heteroaryloxy, amino optionally mono- or di-substituted with alkyl, aryl or heteroaryl groups, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocyclyl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups,
- 25 alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, carboxy, alkoxycarbonyl, aryloxy, heteroaryloxy, or the like attached at any available point to produce a stable compound.

- “Alkynyl” - alone or in combination means a straight or branched hydrocarbon
- 30 containing 2-20, preferably 2-17, more preferably 2-10, even more preferably 2-8, most

WO 01/40246

PCT/US00/32721

preferably 2-4, carbon atoms containing at least one, preferably one, carbon to carbon triple bond. Examples of alkynyl groups include ethynyl, propynyl, butynyl and the like. A substituted alkynyl refers to the straight chain alkynyl or branched alkenyl defined previously, independently substituted with 1 to 3 groups or substituents of halo, hydroxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, acyloxy, aryloxy, heteroaryloxy, amino optionally
 5 mono- or di-substituted with alkyl, aryl or heteroaryl groups, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocyclyl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups, alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, or the like attached at any available point to produce a stable
 10 compound.

“Alkyl alkenyl” refers to a group $-R-CR'=CR''R'''$, where R is lower alkyl, or substituted lower alkyl, R', R'', R''' may independently be hydrogen, halogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as defined
 15 below.

“Alkyl alkynyl” refers to a groups $-RCR'$ where R is lower alkyl or substituted lower alkyl, R' is hydrogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as defined below.

“Alkoxy” denotes the group $-OR$, where R is lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroalkyl, heteroarylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, or substituted cycloheteroalkyl as defined.
 20

“Alkylthio” denotes the group $-SR$, $-S(O)_{n=1,2}-R$, where R is lower alkyl, substituted lower alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl as defined herein.

“Acyl” denotes groups $-C(O)R$, where R is hydrogen, lower alkyl substituted lower alkyl, aryl, substituted aryl and the like as defined herein.
 25

“Aryloxy” denotes groups $-OAr$, where Ar is an aryl, substituted aryl, heteroaryl, or substituted heteroaryl group as defined herein.

“Amino” denotes the group NRR' , where R and R' may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as
 30 defined herein or acyl.

WO 01/40246

PCT/US00/32721

“Amido” denotes the group -C(O)NRR', where R and R' may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, substituted hetaryl as defined herein.

“Carboxyl” denotes the group -C(O)OR, where R is hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, and substituted hetaryl as defined herein.

“Aryl” - alone or in combination means phenyl or naphthyl optionally carbocyclic fused with a cycloalkyl of preferably 5-7, more preferably 5-6, ring members and/or optionally substituted with 1 to 3 groups or substituents of halo, hydroxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, acyloxy, aryloxy, heteroaryloxy, amino optionally mono- or di-substituted with alkyl, aryl or heteroaryl groups, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocyclyl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups, alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, or the like.

“Substituted aryl” refers to aryl optionally substituted with one or more functional groups, *e.g.*, halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Heterocycle” refers to a saturated, unsaturated, or aromatic carbocyclic group having a single ring (*e.g.*, morpholino, pyridyl or furyl) or multiple condensed rings (*e.g.*, naphthpyridyl, quinoxalyl, quinolinyl, indolizinyll or benzo[b]thienyl) and having at least one hetero atom, such as N, O or S, within the ring, which can optionally be unsubstituted or substituted with, *e.g.*, halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Heteroaryl” - alone or in combination means a monocyclic aromatic ring structure containing 5 or 6 ring atoms, or a bicyclic aromatic group having 8 to 10 atoms, containing one or more, preferably 1-4, more preferably 1-3, even more preferably 1-2, heteroatoms independently selected from the group O, S, and N, and optionally substituted with 1 to 3 groups or substituents of halo, hydroxy, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl,

WO 01/40246

PCT/US00/32721

acyloxy, aryloxy, heteroaryloxy, amino optionally mono- or di-substituted with alkyl, aryl or heteroaryl groups, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocyclyl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups, alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, or the like. Heteroaryl is
5 also intended to include oxidized S or N, such as sulfinyl, sulfonyl and N-oxide of a tertiary ring nitrogen. A carbon or nitrogen atom is the point of attachment of the heteroaryl ring structure such that a stable aromatic ring is retained. Examples of heteroaryl groups are pyridinyl, pyridazinyl, pyrazinyl, quinazolinyl, purinyl, indolyl, quinolinyl, pyrimidinyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, isoxazolyl, oxathiadiazolyl, isothiazolyl, tetrazolyl,
10 imidazolyl, triazinyl, furanyl, benzofuryl, indolyl and the like. A substituted heteroaryl contains a substituent attached at an available carbon or nitrogen to produce a stable compound.

“Heterocyclyl” - alone or in combination means a non-aromatic cycloalkyl group having from 5 to 10 atoms in which from 1 to 3 carbon atoms in the ring are replaced by
15 heteroatoms of O, S or N, and are optionally benzo fused or fused heteroaryl of 5-6 ring members and/or are optionally substituted as in the case of cycloalkyl. Heterocyclyl is also intended to include oxidized S or N, such as sulfinyl, sulfonyl and N-oxide of a tertiary ring nitrogen. The point of attachment is at a carbon or nitrogen atom. Examples of heterocyclyl
20 groups are tetrahydrofuranyl, dihydropyridinyl, piperidinyl, pyrrolidinyl, piperazinyl, dihydrobenzofuryl, dihydroindolyl, and the like. A substituted heterocyclyl contains a substituent nitrogen attached at an available carbon or nitrogen to produce a stable compound.

“Substituted heteroaryl” refers to a heterocycle optionally mono or poly substituted with one or more functional groups, e.g., halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, substituted
25 heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Aralkyl” refers to the group -R-Ar where Ar is an aryl group and R is lower alkyl or substituted lower alkyl group. Aryl groups can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl,
30 aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano,

WO 01/40246

PCT/US00/32721

thiol, sulfamido and the like.

“Heteroalkyl” refers to the group -R-Het where Het is a heterocycle group and R is a lower alkyl group. Heteroalkyl groups can optionally be unsubstituted or substituted with *e.g.*, halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Heteroarylalkyl” refers to the group -R-HetAr where HetAr is an heteroaryl group and R lower alkyl or substituted lower alkyl. Heteroarylalkyl groups can optionally be unsubstituted or substituted with, *e.g.*, halogen, lower alkyl, substituted lower alkyl, alkoxy, alkylthio, acetylene, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Cycloalkyl” refers to a divalent cyclic or polycyclic alkyl group containing 3 to 15 carbon atoms.

“Substituted cycloalkyl” refers to a cycloalkyl group comprising one or more substituents with, *e.g.*, halogen, lower alkyl, substituted lower alkyl, alkoxy, alkylthio, acetylene, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Cycloheteroalkyl” refers to a cycloalkyl group wherein one or more of the ring carbon atoms is replaced with a heteroatom (*e.g.*, N, O, S or P).

“Substituted cycloheteroalkyl” refers to a cycloheteroalkyl group as herein defined which contains one or more substituents, such as halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Alkyl cycloalkyl” denotes the group -R-cycloalkyl where cycloalkyl is a cycloalkyl group and R is a lower alkyl or substituted lower alkyl. Cycloalkyl groups can optionally be unsubstituted or substituted with *e.g.* halogen, lower alkyl, lower alkoxy, alkylthio, acetylene, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

“Alkyl cycloheteroalkyl” denotes the group -R-cycloheteroalkyl where R is a lower alkyl or substituted lower alkyl. Cycloheteroalkyl groups can optionally be unsubstituted or

WO 01/40246

PCT/US00/32721


substituted with *e.g.* halogen, lower alkyl, lower alkoxy, alkylthio, amino, amido, carboxyl, acetylene, hydroxyl, aryl, aryloxy, heterocycle, substituted heterocycle, hetaryl, substituted hetaryl, nitro, cyano, thiol, sulfamido and the like.

The compounds of this invention can be prepared as outlined in the schemes 1-5
5 below. A general outline for the preparation of **V** and **VI** is shown in Scheme 1. Compound **I** can be prepared, following the procedures reported earlier (U.S. Patent No. 5,789,416, the specification of which is incorporated herein by reference), by reacting 6-chloropurine riboside **1** with a primary amine R^1NH_2 . The 2', 3' hydroxy groups can be protected as acetonide by reacting **I** with 2,2'-dimethoxypropane in the presence of a catalytic amount of
10 TsOH [Evans, Parrish and Long Carbohydrat. Res., 3, 453 (1967)] to give **II**. Activation of the 5'-hydroxyl of **II** with MsCl in pyridine can give the 5'-mesylate **III**. Displacement of the 5'-mesylate with R^5SNa can give sulfides with the general formula **IV**. Treatment of **IV** with an acid can free the 2', 3' hydroxyl groups to give sulfide derivatives with the general formula **V**. Esterification of **V** can afford 2', 3' diesters with the general formula **VI**.

WO 01/40246

PCT/US00/32721

Scheme 1




WO 01/40246

PCT/US00/32721

- The 2-substituted derivatives with the general formula **XV** can be prepared as shown in Scheme 2. Condensation of 1,2,3,5-tetraacetylribofuranaside **2** with 2-substituted-6-chloropurine **VII** can give 2-substituted-6-chloropurineriboside triacetate **VIII** which on reaction with a primary amine R^1NH_2 can give 2-substituted-6-alkylamino derivatives **IX**.
- 5 Hydrolysis of the acetates followed by protection of the 2', 3' hydroxy groups as an acetonide can give **XI**. Activation of the 5'-hydroxyl of **XI** with $MsCl$ in pyridine can give the 5'-mesylate **XII**. Displacement of the 5'-mesylate with R^5SNa can give sulfides with the general

Scheme 2



formula **XIII** that can be deprotected to give sulfides with general formula **XIV**. Esterification at the 2', 3' positions can afford the 2', 3' diesters with the general formula **XV**.


- 10 Oxidation of sulfides with the general formula **V**, **VI**, **XIV**, **XV** (Scheme 3) with an oxidizing agent (Drabowicz, et.al. The chemistry of sulfones and sulfoxides, Wiley, New

WO 01/40246

PCT/US00/32721

York, 1988, 233-378) can afford corresponding sulfoxides with the general formula XVI, XVII, XVIII, XIX. These sulfoxides on further oxidation can afford sulfones with the general formula XX, XXI, XXII, XXIII.

Scheme 3




- 5 An example of a specific synthesis of one of the compounds of this invention is shown in Scheme 4. Preparation of compound 7 starting from compound 3 is shown in scheme 3. Compound 3 was prepared from 6-chloropurineriboside 1 and 3-(R)-aminotetrahydrofuran following the procedure reported previously (See U.S. Patent No. 5,789,164). Protection of

WO 01/40246

PCT/US00/32721

the 2' and 3' hydroxyls with dimethoxypropane in the presence of TsOH(cat.) gave acetonide 4. Reaction of 4 with MsCl in pyridine at 0 °C gave mesylate 5 which on displacement with sodium methanethiolate in an acetonitrile/water mixture gave sulfide 6. Deprotection of 6 with 80% acetic acid /water gave the target compound 7.


Scheme 4



5

Oxidation of the ethyl sulfide 8 with oxone (Trost, B.M.; Curran, D.P. Tetrahedron Letters 1981, 22, 1287) in MeOH gave sulfone 9 (Scheme 5).

Scheme 5



WO 01/40246

PCT/US00/32721

This invention also includes pro-drugs of the A₁ agonist compositions of this invention. A pro-drug is a drug which has been chemically modified and may be biologically inactive at its site of action, but which will be degraded or modified by one or more enzymatic or *in vivo* processes to the bioactive form. The pro-drugs of this invention should have a different pharmacokinetic profile to the parent enabling improved absorption across the mucosal epithelium, better salt formulation and/or solubility and improved systemic stability. The compounds of this invention may be preferably modified at one or more of the hydroxyl groups to form pro-drugs. The modifications may be (1) ester or carbamate derivatives which may be cleaved by esterases or lipases, for example; (2) peptides which may be recognized by specific or non specific proteinase; or (3) derivatives that accumulate at a site of action through membrane selection or a pro-drug form or modified pro-drug form, or any combination of (1) to (3) above.

If a compound of this invention contains a basic group, then corresponding acid addition salt may be prepared. Acid addition salts of the compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic, or methanesulfonic. The hydrochloric salt form is especially useful. If a compound of this invention contains an acidic group, then corresponding cationic salts may be prepared. Typically the parent compound is treated with an excess of an alkaline reagent, such as hydroxide, carbonate or alkoxide, containing the appropriate cation. Cations such as Na⁺, K⁺, Ca⁺² and NH₄⁺ are examples of cations present in pharmaceutically acceptable salts. Certain of the compounds form inner salts or zwitterions which may also be acceptable.

The compositions of this invention are useful for treating a variety of mammalian disorders and preferably human disorders that are mediated by an A₁ adenosine receptor. For example, the compositions of this invention are useful for modifying cardiac activity in mammals experiencing a coronary electrical disorder that can be treated by stimulating an A₁ adenosine receptor. Examples of coronary electrical disorders that can be treated by the compositions of this invention include supraventricular tachycardias, atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia. Furthermore, orally active A₁ agonists of this invention that demonstrate an excellent safety profile in treating supraventricular arrhythmias

WO 01/40246

PCT/US00/32721

may also be used as a prophylactic for those at high risk of a myocardial ischemia.

The compositions of this invention are also useful for modifying adipocyte function by stimulating an A₁ adenosine receptor that leads to diminished release of NEFA and increased release of leptin. Disease states related to adipocyte function that can be modified using
5 compositions of this invention include diabetes, and obesity.

In skeletal muscle cells, A₁ AdoR agonists mediate a synergistic stimulation of glucose uptake and transport by insulin (Vergauwen, L. *et al*, *J. Clin. Invest.* 1994, 93, 974-81; Challiss, R.A. *et al*, *Eur.J.Pharmacol.*, 1992, 226, 121-8). Another therapeutic utility of compositions of this invention is more efficient regulation of glucose and a decrease of
10 circulating insulin in patients afflicted with diabetes.

The A₁ receptor agonist, R-PIA, has been shown to increase the leptin released from white adipocytes and augment insulin-stimulated leptin production (M. Ozeck Master's Thesis Univ. of Florida 1999 with L. Belardinelli). Evidence suggests that catecholamines inhibit the production of leptin from adipocytes through activation of β -adrenergic receptors.
15 The anti- β -adrenergic effects of A₁ agonists on the adipocytes are believed to play a role in the increased release of leptin. The functional role of leptin is multifaceted including decreased appetite, stimulated energy utilization, and increased fertility.

The compositions of this invention may also be used to provide central nervous system neuroprotection by stimulating an A₁ adenosine receptor. Central nervous system disorders
20 that may be treated using the compositions of this invention include epilepsy, and stroke.

In the kidney, there is evidence that stimulation of the A₁ AdoR promotes sodium retention, promotes exchange of sodium in urine for potassium, and reduces glomerular filtration rate as sodium excretion increases (Gellai, M. *et al*, *JPET*, 1998, 286, 1191-6; Wilcox, C.S. *et al*, *J.Am.Soc.Nephrol.*, 1999, 10, 714-720). It is believed that these responses
25 are elicited by chronic local production of adenosine. That is, in the kidney there is a tonic effect of adenosine to stimulate the A₁ AdoR. Another clinical utility of compositions of this invention, therefore, is the selective antagonism of the A₁ AdoR in the kidney to inhibit sodium retention, inhibit the exchange of sodium for potassium, and preserve kidney glomerular filtration rate when sodium excretion rises to yield a potassium sparing diuretic
30 that preserves renal function.

WO 01/40246

PCT/US00/32721

The compositions of this invention are further useful for providing cardiomyocyte protection from ischemic events by stimulating an A₁ adenosine receptor. Ischemic events treatable using the compositions of this invention include stable angina, unstable angina, cardiac transplant, and myocardial infarction.

5 An important aspect of compounds of this invention is that each compound has an intrinsic efficacy associated with it (for a discussion see T. P. Kenakin Stimulus Response Mechanisms. In Pharmacological Analysis of Drug-Receptor Interaction, Ed. Kenakin, T.P. New York: Raven Press, p 39-68). This intrinsic efficacy is not defined by it's affinity for the receptor, but it is defined as the quantitative effect of the compound to activate a given
10 effector system (eg. cAMP production) in a given cell type. The intrinsic efficacy of a given compound may vary from cell type to cell type and/or from effector system to effector system. When a compound has an intrinsic efficacy lower than a full agonist (i.e. submaximal) than the agonist is called a partial agonist. Thus, a partial agonist is a molecule that binds to a receptor and elicits a response that is smaller than that of a full agonist (submaximal), but also
15 competitively antagonizes the response(s) elicited by a full agonist. The tonic action of adenosine with respect to kidney function is a prime example where a partial A₁ agonist be expected to act as antagonists (e.g. adenosine). The tonic action of adenosine with respect to kidney function is a prime example where a partial A₁ agonist could be expected to act as an antagonist. The compounds of this invention are believed to have therapeutically useful
20 affinities for the adenosine A₁ receptor, and they will have a range of intrinsic efficacies from full agonist to partial agonist. That is, some compounds may have no effect with respect to a given effector system in a given cell type, but be a full agonist in another cell type and/or effector system. The reason for such variable pharmacological behavior relates to the magnitude of the receptor reserve for the A₁ adenosine receptor in any given cell type (eg. AV
25 nodal cells vs. adipocytes) and for a given response. The receptor reserve (spare receptor capacity) is the total number of receptors minus the fraction of receptors that is required to induce the maximal response using a full agonist (L. E. Limbird, Cell Surface Receptors: A Short Course on Theory and Methods, Kluwer Acad. Pub. 1996, Boston, Mass.). Therefore, the agonist could be a full agonist at eliciting a response, and a partial agonist for eliciting
30 another response in other tissue or cells and still be an antagonist or lack activity for a third

WO 01/40246

PCT/US00/32721

response in another tissue or cell. Consequently, a partial agonist targeted to a selected target is likely to cause fewer side effects than a full agonist. As a corollary, a full agonist elicits all the effects mediated by the respective receptor, whereas this is not necessarily the case of a partial agonist. The compounds of this invention based on their affinity for the A₁ receptor and
5 their potency and selectivity to elicit A₁ receptor mediated responses have the potential for therapeutic intervention in the multiple disease states described above.

Partial A₁ agonists may have an added benefit for chronic therapy because they will be less likely to induce desensitization of the A₁ receptor (R. B. Clark, B. J. Knoll, R. Barber TiPS, Vol. 20 (1999) p. 279-286) and to cause side effects. Chronic administration of a full
10 agonist (R-N6-phenylisopropyladenosine, R-PIA) for 7 days led to a desensitization of the A₁ receptor in terms of the dromotropic response in guinea pigs (note: a decrease in receptor number was observed – D. M. Dennis, J. C. Shryock, L. Belardinelli JPET, Vol. 272 (1995) p. 1024-1035). The A₁ agonist induced inhibitory effect on the production of cAMP by adenylate cyclase in adipocytes has been shown to desensitize upon chronic treatment with an
15 A₁ agonist as well (W. J. Parsons and G. L. Stiles J. Biol. Chem. Vol. 262 (1987) p. 841-847).

The compositions of this invention may be administered orally, intravenously, through the epidermis, bolus, nasally, by inhalation or by any other means known in the art for administering a therapeutic agents. The method of treatment comprises the administration of an effective quantity of the chosen compound, preferably dispersed in a pharmaceutical
20 carrier. Dosage units of the active ingredient are generally selected from the range of 0.01 to 100 mg/kg, but will be readily determined by one skilled in the art depending upon the route of administration, age and condition of the patient.

Pharmaceutical compositions including the compounds of this invention, and/or derivatives thereof, may be formulated as solutions or lyophilized powders for parenteral
25 administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. If used in liquid form the compositions of this invention are preferably incorporated into a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water and buffered sodium or ammonium acetate solution. Such liquid formulations are
30 suitable for parenteral administration, but may also be used for oral administration. It may be

WO 01/40246


PCT/US00/32721

desirable to add excipients such as polyvinylpyrrolidinone, gelatin, hydroxycellulose, acacia, polyethylene glycol, mannitol, sodium chloride, sodium citrate or any other excipient known to one of skill in the art to pharmaceutical compositions including compounds of this invention. Alternatively, the pharmaceutical compounds may be encapsulated, tableted or
5 prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Liquid carriers include syrup, peanut oil, olive oil, glycerin, saline, alcohols and water. Solid carriers include starch, lactose, calcium sulfate, dihydrate, teffa alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. The carrier
10 may also include a sustained release material such as glycerol monostearate or glycerol distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 gram per dosage unit. The pharmaceutical dosages are made using conventional techniques such as milling, mixing, granulation, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms.
15 When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly or filled into a soft gelatin capsule.

The Examples which follow serve to illustrate this invention. The Examples are not
20 intended to limit the scope of this invention, but are provided to show how to make and use the compounds of this invention.


WO 01/40246

PCT/US00/32721

Example 1

Intermediate – (4-{6-[(3R)oxolan-3-yl]amino}purin-9-yl)(1R, 2R, 5R)-7,7-dimethyl-3,6,8-trioxabicyclo[3.3.0]oct-2-ylmethan-1-ol (4)

- 5 To a solution of compound 3 (2.0 g, 6.0 mmol) and 2,2-dimethoxypropane (1.2 g, 11.8 mmol) in dimethylformamide (20 mL) was added p-toluenesulfonic acid (50 mg, 0.26 mmol) at 70°C. After 48 h at 70°C, the reaction was concentrated in vacuo to afford a solid. The solid was dissolved in methanol (3 mL), then triturated with ethyl ether (50 mL). The resultant crystals were collected by vacuum filtration to afford the intermediate 4.




10

To a solution of 4 (190 mg, 0.5 mmol) in anhydrous pyridine (5mL), was added MsCl (80 microL, 1 mmol) at 0°C. The reaction mixture was stirred at the same temperature for 2h. Pyridine was removed under reduced pressure, residue was taken in dichloromethane (50mL),

WO 01/40246

PCT/US00/32721


washed with water (3 x 20mL) and dried (Na_2SO_4). Evaporation of the solvent gave product 5 as a white foam: ^1H NMR (CDCl_3) δ 1.4 (s, 3H), 1.6 (s, 3H), 2.0-2.2 (m, 1H), 2.3-2.5 (m, 1H), 2.9 (s, 3H), 3.7-4.2 (m, 4H), 4.4-4.6 (m, 3H), 4.8-5.0 (bs, 1H), 5.1-5.2 (bs, 1H), 5.4-5.5 (bs, 1H), 6.1 (s, 1H), 6.4-6.6 (bs, 1H), 8.1 (s, 1H), 8.4 (s, 1H)



5

A mixture of mesylate 5 (150 mg) and methanethiolate (150mg) in acetonitrile (2mL) and water (1mL) was heated at 70 C for 24h. The solvent was evaporated under reduced pressure and the residue was purified by preparative TLC [methanol-dichloromethane (1:19)] to afford product 6: ^1H NMR (CDCl_3) δ 1.35 (s, 3H), 1.60 (s, 3H), 1.90-2.05 (m, 1H), 2.05 (s, 3H), 2.30-2.40 (m, 1H), 2.70 (doublet of AB quartet, 2H), 3.75-3.90 (m, 2H), 3.95-4.00 (m, 2H), 4.3-4.4 (m, 1H), 4.8-4.95 (m, 1H), 5.00-5.05 (m, 1H), 5.45-5.50 (d, 1H), 6.00-6.10 (m, 2H), 7.85 (s, 1H), 8.3 (s, 1H).

10




2-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(methylthiomethyl)oxolane-3,4-diol (7)

WO 01/40246

PCT/US00/32721


Compound 6 (50mg) was dissolved in a mixture of acetic acid (8 mL) and water (2 mL) and heated at 90°C for 16 h. Solvents were removed under reduced pressure, and the residue was purified by preparative TLC [methanol-dichloromethane (1:9)] to afford compound 7: ¹H NMR (CDCl₃) δ 1.90-2.05 (m, 1H), 2.15 (s, 3H), 2.30-2.40 (m, 1H), 2.75-2.85 (m, 2H), 3.80-3.90 (m, 2H), 3.90-4.00 (m, 2H), 4.30-4.45 (m, 2H), 4.50-4.55 (m, 1H), 4.75-4.95 (m, 1H), 5.90-5.95 (m, 1H), 6.30-6.60 (m, 1H), 7.95 (s, 1H), 8.25 (s, 1H).



8

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(4S,5S,2R,3R)-5-[(Ethylthio)methyl]oxolane-3,4-diol(8)

Compound 8 was prepared in the manner similar to that of 7 substituting ethane thiolate for methane thiolate. (M+1) = 382.30




10

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(4S,5S,2R,3R)-5-[(Methylethylthio)methyl]oxolane-3,4-diol(10)

WO 01/40246

PCT/US00/32721


Compound 10 was prepared in the manner similar to that of 7 substituting i-propane thiolate for methane thiolate. ¹H NMR (CDCl₃) δ 1.25 (d, 6H), 1.90-2.05 (m, 1H), 2.15 (s, 3H), 2.30-2.40 (m, 1H), 2.85-2.87 (d, 2H), 2.95 (septet, 1H), 3.80-3.90 (m, 2H), 3.95-4.05 (m, 2H), 4.35-4.40 (m, 2H), 4.50-4.55 (m, 1H), 4.75-4.85 (m, 1H), 5.90-5.95 (d, 1H), 6.85-6.95 (m, 1H), 7.95 (s, 1H), 8.25 (s, 1H).



11

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(4S,5S,2R,3R)-5-(phenylthiomethyl)oxolane-3,4-diol(11)

Compound 11 was prepared in the manner similar to that of 7 substituting phenyl thiolate for methane thiolate. ¹H NMR (CDCl₃) 1.95-2.05 (m, 1H), 2.30-2.40 (m, 1H), 3.2 (d, 2H), 3.80-3.90 (m, 2H), 3.95-4.10 (m, 2H), 4.35-4.40 (d, 1H), 4.45 (t, 1H), 4.50-4.55 (m, 1H), 4.80-4.90 (m, 1H), 5.85 (d, 1H), 6.70-6.80 (m, 1H), 7.15-7.30 (m, 3H), 7.35 (d, 2H), 7.75 (s, 1H), 8.25 (s, 1H).




12

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(4S,5S,2R,3R)-5-[(4-Methoxyphenylthio)methyl]oxolane-3,4-diol(12)

WO 01/40246


PCT/US00/32721

This compound was prepared in the manner similar to that of 7 substituting 4-methoxyphenyl thiolate for methane thiolate. (M+1) = 460.4



- 5 **2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}(4S,5S,2R,3R)-5-[(4-chlorophenylthio)methyl]oxolane-3,4-diol(13)**

This compound was prepared in a manner similar to that of 7 substituting 4-chlorophenyl thiolate for methane thiolate. (M+1) = 464.3




- 10 **2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}(4S,5S,2R,3R)-5-[(4-fluorophenylthio)methyl]oxolane-3,4-diol(14)**

This compound was prepared in a manner similar to that of 7 substituting 4-fluorophenyl thiolate for methane thiolate. (M+1) = 448.3


WO 01/40246

PCT/US00/32721

**15**


2-{6-[(3R)oxolan-3-ylamino]purin-9-yl}(4S,5S,2R,3R)-5-[(4-methylphenylthio)methyl]oxolane-3,4-diol(15)

This compound was prepared in a manner similar to that of 7 substituting 4-methylphenyl thiolate for methane thiolate. (M+1) = 444.38

**16**

2-{6-[(3R)oxolan-3-ylamino]purin-9-yl}(4S,5S,2R,3R)-5-[(4-(trifluoromethyl)phenylthio)methyl]oxolane-3,4-diol(16)

This compound was prepared in a manner similar to that of 7 substituting 4-trifluoromethylphenyl thiolate for methane thiolate. (M+1) = 488.36


**17**

2-{6-[(3R)oxolan-3-ylamino]purin-9-yl}(4S,5S,2R,3R)-5-[(2-Methoxyphenylthio)methyl]oxolane-3,4-diol(17)

WO 01/40246

PCT/US00/32721

This compound was prepared in a manner similar to that of 7 substituting 2-methoxyphenyl thiolate for methane thiolate. (M+1) = 460.4



9


(5-{6-[(3R)oxolan-3-yl]amino}purin-9-yl){(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl}(ethylsulfonyl)methane(9)

5

To a cooled solution of sulfide 8 in methanol at 0°C under nitrogen was added 3 eq. of Oxone (Potassium peroxy monosulfate) and the reaction mixture was allowed to stir at the same temperature for 1 hour. After the starting material consumed (by TLC), the reaction mixture was concentrated and filtered through a small plug of silica gel. Purification by preparative

10 TLC [methanol-dichloromethane (1:19)] afforded 9 as an off-white hygroscopic solid.

(M+1) = 414.28



18


2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl){(4S,5S,2R,3R)-5-[(2,4-difluorophenylthio)methyl]oxolane-3,4-diol(18)

15

This compound was prepared in a manner similar to that of 7 substituting 2,4-difluorophenyl thiolate for methane thiolate. (M+1) = 466.23

WO 01/40246

PCT/US00/32721




19

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}(4S,5S,2R,3R)-5-[(2,6-dichlorophenylthio)methyl]oxolane-3,4-diol(19)

This compound was prepared in a manner similar to that of 7 substituting 2,6-dichlorophenyl thiolate for methane thiolate. (M+1) = 498.18

5



20


2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}(4S,5S,2R,3R)-5-[(3-fluorophenylthio)methyl]oxolane-3,4-diol(20)

This compound was prepared in a manner similar to that of 7 substituting 3-fluorophenyl thiolate for methane thiolate. (M+1) = 448.26

10

WO 01/40246


PCT/US00/32721



21

2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(4S,5S,2R,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol(21)

This compound was prepared in a manner similar to that of 7 substituting 2-fluorophenyl thiolate for methane thiolate. (M+1) = 448.24




22

5-{6-[(3R)oxolan-3-yl]amino}purin-9-yl-(2S,3R,4R,5R)-4-acetyloxy-2-[(fluorophenylthio)methyl]oxolan-3-yl acetate(22)

To a solution of compound 21 (139 mg) in pyridine (2 mL) at 23 °C was added acetic anhydride (0.1 mL). After 3 h at 23 °C, the reaction was concentrated *in vacuo*. The residue was dissolved in methylene chloride (50 mL), washed with water (3 x 10 mL), and dried (Na₂SO₄). After concentration *in vacuo*, the residue was purified by flash chromatography (methylene chloride: methanol 20:1 followed by 9:1) to afford compound 22 (170 mg):

WO 01/40246


PCT/US00/32721



23

Methyl 2[(5-{6-(((3R)oxolan-3-yl)amino)purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]benzoate (23)

To a solution of Compound 4 (0.377g, 1 mmol) in 5mL of THF, was added Triphenylphosphine (0.524g, 2 mmol), DEAD (0.40 mL, 2 mmoles), let stir for 5 minutes before adding 2-carbomethoxythiophenol (0.5mL). Reaction was allowed to stir under reflux. After 72 h of reflux, the reaction was concentrated in vacuo and the residue purified by flash column chromatography (20%EtOAc/Hexanes) to give a clear viscous oil. It was taken into a mixture of acetic acid (8mL) and water (2mL) and heated at 80 C for 16h. Solvents were removed in vacuo and the residue was purified by prep TLC [methanol-dichloromethane (1:9)] to give compound 23. (M+1) = 488.5




24

{2[(5-{6-(((3R)oxolan-3-yl)amino)purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]phenyl}-N-methylcarboxamidebenzoate (24)

Compound 23 was taken into 40% aq.methylamine (2 mL) and I-propanol (2 mL) and heated at 70 C for 16h. Solvents were removed in vacuo and the residue was purified by prep TLC TLC [methanol-dichloromethane (1:9)] to give compound 24. (M+1) = 487.5

WO 01/40246


PCT/US00/32721



25

2-{6-[(3R)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(benzoxazol-2-ylthiomethyl)oxolane-3,4-diol (25)

This compound was prepared in a manner similar to that of 23 substituting 2-mercaptobenzoxazole for 2-carbomethoxy thiophenol ($M+1$) = 471.4




26

5 2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-[(1-methylimidazol-2-yl-thio)methyl]oxolane-3,4-diol (26)

Compound 26 was prepared in the manner of compound 23 substituting 2-mercapto-1-methylimidazole for 2-carbomethoxythiophenol [MS 434.4 ($M+1$)].

2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(pyrimidine-2-




27

WO 01/40246

PCT/US00/32721


ylthiomethyl)oxolane-3,4-diol (27)

Compound 27 was prepared in the manner of compound 23 substituting 2-mercaptopyrimidine for 2-carbomethoxythiophenol [MS 432.4 (M+1)].



5 **2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(2-pyridylthiomethyl)oxolane-3,4-diol (28)**

Compound 28 was prepared in the manner of compound 23 substituting 2-mercaptopyridine for 2-carbomethoxythiophenol [MS 431.4 (M+1)].




10 **2-{6-[(3S)oxolan-3-yl]amino]purin-9-yl}(4S,5S,2R,3R)-5-(4-pyridylthiomethyl)oxolane-3,4-diol (29)**

Compound 29 was prepared in the manner of compound 23 substituting 4-mercaptopyridine for 2-carbomethoxythiophenol [MS 431.4 (M+1)].

WO 01/40246

PCT/US00/32721



5- $\{6-[(3R)oxolan-3-yl]amino\}purin-9-yl\}(2S,3R,4R,5R)-4-acetyloxy-2-[(4-fluorophenylthio)methyl]oxolan-3-yl\}acetate$ (30) (M+1) = 532.17.

5

EXAMPLE 2

Binding Assays – DDT₁ Cells

Cell Culture

DDT cells (hamster vas deferens smooth muscle cell line) were grown as monolayers in petri dishes using Dulbecco's Modified Eagle's Medium (DMEM) containing 2.5 g ml⁻¹ amphotericin B, 100 U ml⁻¹ penicillin G, 0.1 mg ml⁻¹ streptomycin sulfate and 5% fetal bovine serum in a humidified atmosphere of 95% air and 5% CO₂. Cells were subcultured twice weekly by dispersion in Hank's Balanced Salt Solution (HBSS) without the divalent cations and containing 1 mM EDTA. The cells were then seeded in growth medium at a density of 1.2 x 10⁵ cells per plate and experiments were performed 4 days later at approximately one day preconfluence.

15

Membrane Preparations

Attached cells were washed twice with HBSS (2 x 10 ml), scraped free of the plate with the aid of a rubber policeman in 5 ml of 50 mM Tris-HCl buffer pH 7.4 at 4 °C and the suspension homogenized for 10 s. The suspension was then centrifuged at 27,000 x g for 10 min. The pellet was resuspended in homogenization buffer by vortexing and centrifuged as described above. The final pellet was resuspended in 1 vol of 50 mM Tris-HCl buffer pH 7.4 containing 5 mM MgCl₂ for A₁ AdoR assays. For the [³⁵S]GTPγS binding assay the final

20

WO 01/40246

PCT/US00/32721

pellet was resuspended in 50 mM Tris-HCl pH 7.4 containing 5 mM MgCl₂, 100 mM NaCl and 1 mM dithiothreitol. This membrane suspension was then placed in liquid nitrogen for 10 min, thawed and used for assays. The protein content was determined with a Bradford™ Assay Kit using bovine serum albumin as standard.

5 Competitive Binding Assay

Pig striatum were prepared by homogenation in 50 mM Tris buffer (5x volume of tissue mass pH = 7.4). After centrifugation at 19,000 rpm for 25 minutes at 4 °C, the supernatant was discarded, and the process was repeated twice. Compositions of this invention were assayed to determine their affinity for the A₁ receptor in a pig striatum membrane prep or a DDT₁ membrane prep. Briefly, 0.2 mg of pig striatal membranes or DDT₁ cell membranes were treated with adenosine deaminase and 50 mM Tris buffer (pH = 7.4) followed by mixing. To the pig membranes was added 2 µL of serially diluted DMSO stock solution of the compounds of this invention at concentrations ranging from 100 microM to 10 nM. The control received 2 microL of DMSO alone, then the antagonist [³H] 8-cyclopentylxanthine (CPX) for pig striatum or the agonist [³H] 2-chloro-6-cyclopentyladenosine (CCPA) for DDT₁ membranes in Tris buffer (50 mM, pH of 7.4) was added to achieve a final concentration of 2 nM. After incubation at 23 C for 2h, then the solutions were filtered using a membrane harvester using multiple washing of the membranes (3 x). The filter disks were counted in scintillation cocktail affording the amount of displacement of tritiated CPX or by the competitive binding compositions of this invention. Greater than a 5 point curve was used to generate K_i's and the number of experiments is indicated in the column marked in Table 1, below:

Table 1

Compound #	K _i – DDT ₁ cell membrane	K _i – Pig Striatum
7	--	222 nM
10	--	188 nM
11	--	44 nM
12	820 nM	--
14	363 nM	--
15	922 nM	--
16	7701 nM	--
17	947 nM	--

WO 01/40246

PCT/US00/32721

EXAMPLE 3**[³⁵S]GTPγS Binding Assays**

A₁-agonist stimulated [³⁵S] GTPγS binding was determined by a modification of the method described by Gierschik et al. (1991) and Lorenzen et al. (1993). Membrane protein (30-50 μg) was incubated in a volume of 0.1 ml containing 50 mM Tris-HCl buffer pH 7.4, 5 mM MgCl₂, 100 mM NaCl, 1 mM dithiothreitol, 0.2 units ml⁻¹ adenosine deaminase, 0.5% BSA, 1 mM EDTA, 10 mM GDP, 0.3 nM [³⁵S]GTPγS and with or without varying concentrations of CPA for 90 min at 30 °C. Nonspecific binding was determined by the addition of 10 μM GTPγS. Agonist stimulated binding was determined as the difference between total binding in the presence of CPA and basal binding determined in the absence of CPA. Previous reports have shown that agonist stimulated [³⁵S]GTPγS binding was dependent on the presence of GDP (Gierschik et al., 1991; Lorenzen et al., 1993; Traynor & Nahorski, 1995). In preliminary experiments, it was found that 10 μM GDP gave the optimal stimulation of CPA dependent [³⁵S]GTPγS binding and this concentration was therefore used in all studies. In saturation experiments, 0.5 nM [³⁵S]GTPγS was incubated with 0.5-1000 nM GTPγS. At the end of the incubation, each suspension was filtered and the retained radioactivity determined as described above. Results are presented normalized to the full agonist N-6-cyclopentyladenosine, CPA.

Table 2

Compound #	GTPγS
CPA	100 %
8	104%
12	52%
13	69%
14	61%
15	48%
16	31%
17	52%

WO 01/40246

PCT/US00/32721

EXAMPLE 4**cAMP Assay**

A scintillation proximity assay (SPA) using rabbit antibodies directed at cAMP using an added tracer of adenosine 3',5'-cyclic phosphoric acid 2'-O-succinyl-3-[¹²⁵I]iodotyrosine methyl ester and fluoromicrospheres containing anti-rabbit specific antibodies as described by Amersham Pharmacia Biotech (Biotrak cellular communication assays). Briefly, DDT₁ cells were cultured in clear bottomed 96 well microtiter plates with opaque wells at concentrations between 10⁴ to 10⁶ cells per well in 40 µl of HBSS at 37 °C (5% CO₂ and 95% humidity). The partial or full A₁ agonists (5 µl) of this invention were incubated at various concentrations with the DDT₁ cells in the presence of rolipram (50 µM), and 5 µM forskolin for 10 min at 37 °C. The cells were immediately lysed by treatment 5 µl of 10% dodecyltrimethylammonium bromide followed by shaking using microplate shaker. After incubation of the plate for 5 minutes, an immunoreagent solution (150 µl containing equal volumes of tracer, antiserum, and SPA fluorospheres) was added to each well followed by sealing the plate. After 15-20 h at 23 °C, the amount of bound [¹²⁵I] cAMP to the fluoromicrospheres was determined by counting in a microtitre plate scintillation counter for 2 minutes. Comparison of counts with standard curves generated for cAMP using a similar protocol afforded the cAMP present after cell lysis. Results are presented normalized to the full agonist N-6-cyclopentyladenosine, CPA. Thus, the full agonist CPA diminished the amount of forskolin induced cAMP generation back to basal levels.

Table 3


Compound #	Camp
CPA	107 %
8	37%
12	-9%
13	30%
14	47%
15	22%
16	22%
17	18%

WO 01/40246

PCT/US00/32721

What we claim is:

1. A composition of matter having the formula:



5 wherein $X' = S, S(O), S(O)_2$;

R^1 is a monocyclic or polycyclic heterocyclic group containing from 3 to 15 carbon atoms wherein at least one carbon atom is substituted with an atom or molecule selected from the group consisting of N, O, P and $S-(O)_{0,2}$ wherein R_1 does not contain an epoxide group;

R_2 is selected from the group consisting of hydrogen, halo, CF_3 , and cyano;

10 R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(CO)-R'$, and $-(CO)-R''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CO_2R^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $N(R^{20})_2$, $NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{20}C(NR^{20})NHR^{23}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $OCONR^{20}SO_2R^{22}$, $OC(O)R^{20}$, $C(O)OCH_2OC(O)R^{20}$, and $OCON(R^{20})_2$ and wherein each optional heteroaryl, aryl, and heterocyclyl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, $NR^{20}COR^{22}$, $NR^{20}SO_2R^{22}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $NR^{20}CON(R^{20})_2$,
20

WO 01/40246

PCT/US00/32721

OC(O)R²⁰, OC(O)N(R²⁰)₂, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, CN, or OR²⁰;

R₅ is selected from the group consisting of C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, wherein each alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, NO₂, heterocyclyl, aryl, heteroaryl, CF₃, CN, OR²⁰, SR²⁰, S(O)₃R²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, SO₂NR²⁰COR²², SO₂NR²⁰CO₂R²², SO₂NR²⁰CON(R²⁰)₂, P(O)(OR²⁰)₂, N(R²⁰)₂, NR²⁰COR²², NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, NR²⁰C(NR²⁰)NHR²³, COR²⁰, CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²², OC(O)R²⁰, C(O)OCH₂OC(O)R²⁰, and OCON(R²⁰)₂ and wherein the optional heteroaryl, aryl, and heterocyclyl substituent are each further optionally substituted with halo, NO₂, alkyl, CF₃, amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, NR²⁰COR²², NR²⁰SO₂R²², COR²⁰, CO₂R²⁰, CON(R²⁰)₂, NR²⁰CON(R²⁰)₂, OC(O)R²⁰, OC(O)N(R²⁰)₂, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, CN, or OR²⁰;

R²⁰ is selected from the group consisting of H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN, O-C₁₋₆ alkyl, CF₃, aryl, and heteroaryl; and

R²² is selected from the group consisting of C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, which alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN, O-C₁₋₆ alkyl, CF₃, and heteroaryl.

2. The composition of claim 1 wherein R₂ is selected from the group consisting of hydrogen, and halo;

R₃ and R₄ are each independently selected from the group consisting of hydrogen, –(CO)-R', and –(CO)-R'' wherein R' and R'' are each independently selected from the group consisting of C₁₋₁₅ alkyl, heterocyclyl, aryl, and heteroaryl, which alkyl, aryl, heterocyclyl, and heteroaryl are each optionally substituted with from 1 to 2 substituents independently selected from the group of halo, NO₂, heterocyclyl, aryl, heteroaryl, CF₃, CN, OR²⁰, S(O)R²², SO₂R²²,

WO 01/40246

PCT/US00/32721

$\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{C}(\text{NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$ and wherein each optional heteroaryl, aryl, and heterocyclyl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, CN, or OR^{20} ;

- 5 R_5 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group of halo, alkyl, heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, $\text{S}(\text{O})_3\text{R}^{20}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, and
- 10 wherein each optional heteroaryl, and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, SR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN, or OR^{20} ;

- R^{20} is selected from the group consisting of H, C_{1-15} alkyl, aryl, and heteroaryl, which alkyl, aryl, and heteroaryl are each optionally substituted with from 1 to 2 substituents
- 15 independently selected from halo, alkyl, mono- or dialkylamino, CN, $\text{O}-\text{C}_{1-6}$ alkyl, CF_3 ; and

R^{22} is selected from the group consisting of C_{1-15} alkyl, aryl, and heteroaryl, which alkyl, aryl, and heteroaryl are each optionally substituted with from 1 to 2 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or CN, $\text{O}-\text{C}_{1-6}$ alkyl, and CF_3 .

- 20 3. The composition of claim 1 wherein R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-10} alkyl, aryl, and heteroaryl, which alkyl, aryl, and heteroaryl are optionally substituted with from 1 to 2 substituents independently selected from the group of halo, NO_2 , aryl, heteroaryl, CF_3 , CN, OR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{NR}^{20}\text{COR}^{22}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, and wherein each optional heteroaryl, aryl, and heterocyclyl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 ;

25

R_5 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, heterocyclyl, aryl, and heteroaryl, wherein each alkyl, alkenyl, aryl, heterocyclyl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of

30

WO 01/40246

PCT/US00/32721

halo, alkyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN , and OR^{20} ;

5 R^{20} is selected from the group consisting of H, C_{1-6} alkyl, and aryl, which alkyl, and aryl, are optionally substituted with 1 substituent selected from halo, alkyl, mono- or dialkylamino, CN , O-C_{1-6} alkyl, CF_3 ; and

R^{22} is selected from the group consisting of C_{1-6} alkyl and aryl, which alkyl and aryl are optionally substituted with 1 substituent independently selected from halo, alkyl, mono- or dialkylamino, alkyl or CN , O-C_{1-6} alkyl, and CF_3 .

4. The composition of claim 1 wherein R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl, and aryl, which alkyl and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group of halo, NO_2 , aryl, CF_3 , CN , OR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{N}(\text{R}^{20})_2$, and wherein each optional aryl substituent is optionally substituted with halo, NO_2 , alkyl, CF_3 ;

R_5 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkenyl, aryl, and heteroaryl, wherein alkyl, alkenyl, aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN , or OR^{20} ;

25 R^{20} is selected from the group consisting of H, C_{1-6} alkyl, and aryl, which alkyl and aryl are optionally substituted with 1 substituent selected from halo, alkyl, mono- or dialkylamino, CN , O-C_{1-6} alkyl, CF_3 ; and

R^{22} is selected from the group consisting of C_{1-6} alkyl and aryl, which alkyl and aryl are optionally substituted with 1 substituent selected from halo, alkyl or CN , O-C_{1-6} alkyl, and CF_3 .

WO 01/40246

PCT/US00/32721

5. The composition of claim 1 wherein R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(CO)-R'$ and $-(CO)-R''$ wherein each R' and R'' are independently selected from the group consisting of C_{1-6} alkyl which alkyl are optionally substituted with 1 substituent selected from the group of aryl, CF_3 , CN, OR^{20} , $N(R^{20})_2$, and wherein each optional aryl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 ;

R_5 is selected from the group consisting of C_{1-8} alkyl, C_{2-8} alkenyl, and aryl wherein alkyl, alkenyl, and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $NR^{20}CON(R^{20})_2$, CO_2R^{20} , $CON(R^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN, and OR^{20} ;

R^{20} is selected from the group consisting of H, C_{1-6} alkyl; and

R^{22} is selected from the group consisting of C_{1-6} .

6. The composition of claim 1 wherein $X^1=S$ or SO_2 ;

R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(CO)-R'$ and $-(CO)-R''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl;

R_5 is selected from the group consisting of C_{1-8} alkyl, and aryl wherein alkyl, and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $NR^{20}CON(R^{20})_2$, CO_2R^{20} , $CON(R^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN, and OR^{20} ;

R^{20} is selected from the group consisting of H, C_{1-6} alkyl; and

R^{22} is selected from the group consisting of C_{1-6} .

7. The composition of claim 1 wherein $X^1=S$ or SO_2 ;

R_2 is a hydrogen;

R_3 and R_4 are hydrogen;

R_5 is selected from the group consisting of C_{1-8} alkyl, and aryl wherein alkyl, and aryl

WO 01/40246

PCT/US00/32721

are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, CF_3 , CN , OR^{20} , CO_2R^{20} , and

R^{20} is selected from the group consisting of H, C_{1-6} alkyl.

8. The composition of claim 1 wherein $\text{X}^1=\text{S}$ or SO_2 ;

5 R_2 is a hydrogen;

R_3 and R_4 are hydrogen;

R_5 is selected from the group consisting of C_{1-8} alkyl, and aryl wherein alkyl, and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, CF_3 , OR^{20} ; and

10 R^{20} is selected from the group consisting of H, C_{1-6} alkyl.

9. The composition of claim 1 wherein $\text{X}^1=\text{S}$ or SO_2 ;

R_2 is a hydrogen;

R_3 and R^4 are independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl which alkyl are optionally substituted with 1 substituent selected from the group consisting of aryl, CF_3 , CN , OR^{20} , $\text{N}(\text{R}^{20})_2$, and wherein each optional aryl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 ;

R_5 is C_{1-8} alkyl, wherein alkyl, is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN , and OR^{20} ;

R^{20} is selected from the group consisting of H, C_{1-6} alkyl; and

R^{22} is selected from the group consisting of C_{1-6} .

25 10. The composition of claim 1 wherein $\text{X}^1=\text{S}$ or SO_2 ;

R_2 is a hydrogen;

R_3 and R_4 are independently selected from the group consisting of hydrogen, $-(\text{CO})-\text{R}'$ and $-(\text{CO})-\text{R}''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl;

30 R_5 is C_{1-8} alkyl that is optionally substituted with from 1 to 2 substituents

WO 01/40246

PCT/US00/32721

independently selected from the group consisting of aryl, heteroaryl, OR^{20} , $S(O)R^{22}$, CO_2R^{20} , $CON(R^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , CN , and OR^{20} ;

R^{20} is selected from the group consisting of H, C_{1-3} alkyl; and

5 R^{22} is selected from the group consisting of C_{1-6} .

11. The composition of claim 1 wherein $=S$ or SO_2 ;

R_2 is a hydrogen;

R_3 and R_4 are hydrogen;

10 R_5 is C_{1-8} alkyl that is optionally substituted with 1 substituent selected from the group consisting of CO_2R^{20} , and $CON(R^{20})_2$; and

R^{20} is selected from the group consisting of H, and methyl.

12. The composition of claim 11 wherein R_5 is C_{1-6} alkyl.

13. The composition of claim 11 wherein R_5 is selected from the group consisting of methyl and ethyl and isopropyl.

15 14. The composition of claim 1 wherein R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(CO)-R'$ and $-(CO)-R''$ wherein each R' and R'' are independently selected from the group consisting of C_{1-6} alkyl, and aryl, which alkyl and aryl are optionally substituted with from 1 to 2 substituents independently selected from the group of halo, NO_2 , aryl, CF_3 , CN , OR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $N(R^{20})_2$, and wherein each optional aryl substituent is further optionally substituted with halo, NO_2 , alkyl, CF_3 ;

25 R_5 is selected from the group consisting of, aryl, and heteroaryl, wherein aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, CO_2R^{20} , $CON(R^{20})_2$, and wherein each optional heteroaryl, and aryl substituent is further optionally substituted with halo, alkyl, CF_3 , CO_2R^{20} , $CON(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, CN , or OR^{20} ;

30 R^{20} is selected from the group consisting of H, C_{1-6} alkyl, and aryl, which alkyl and aryl are optionally substituted with 1 substituent selected from halo, alkyl, mono- or dialkylamino, CN , $O-C_{1-6}$ alkyl, CF_3 ; and

WO 01/40246

PCT/US00/32721

R^{22} is selected from the group consisting of C_{1-6} alkyl and aryl, which alkyl and aryl are optionally substituted with 1 substituent selected from halo, alkyl or CN, $O-C_{1-6}$ alkyl, and CF_3 .

15. The composition of claim 1 wherein $X^1=S$;

5 R_2 is a hydrogen;

R_3 and R_4 are each independently selected from the group consisting of hydrogen, $-(CO)-R'$ and $-(CO)-R''$ wherein R' and R'' are each independently selected from the group consisting of C_{1-6} alkyl;

10 R_5 is selected from the group consisting of, aryl, and heteroaryl, wherein aryl, and heteroaryl are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF_3 , CN, OR^{20} , SR^{20} , CO_2R^{20} , $CON(R^{20})_2$; and

R^{20} is selected from the group consisting of H, C_{1-3} alkyl.

16. The composition of claim 1 wherein $X^1=S$;

R_2 is a hydrogen;

15 R_3 and R_4 are hydrogen;

R_5 is aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, alkyl, CF_3 , OR^{20} , CO_2R^{20} , $CON(R^{20})_2$;

R^{20} is selected from the group consisting of H, and methyl; and

R^{22} is selected from the group consisting of C_{1-6} alkyl.

20 17. The composition of claim 16 wherein R_5 is phenyl that is optionally substituted with a substituent selected from the group consisting of methoxy, chloro, fluoro, methyl, and trifluoromethyl.

18. The composition of claim 1 wherein R^1 is mono or polysubstituted with one or more compounds selected from the group consisting of halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof.

19. The composition of matter of claim 1 wherein R^1 is a monocyclic, bicyclic, or tricyclic cycloalkyl group containing from 3 to 15 carbon atoms wherein at least one carbon atom is substituted with an atom or molecule selected from the group consisting of O or S-

30


WO 01/40246

PCT/US00/32721

(O)₀₋₂.

20. The composition of claim 19 wherein R¹ is mono or polysubstituted with one or more compounds selected from the group consisting of halogen, oxo, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof.

21. The composition of claim 1 wherein R¹ is:



- 10 wherein R₁['], R₁^{''}, R₁^{'''}, and R₁^{''''} are each independently selected from the group halogen, hydroxyl, lower alkyl, substituted lower alkyl, alkoxy, aryl, acyl, aryloxy, carboxyl, substituted aryl, heterocycle, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, nitro, cyano and mixtures thereof and X is O, or S (-O)₀₋₂.

- 15 22. The composition of claim 21 wherein R₁^{'''} and R₁^{''''} can together be a single oxygen atom.

23. The composition of claim 21 wherein R₁['], R₁^{''}, R₁^{'''}, and R₁^{''''} are each individually selected from the group H, lower alkyl, substitute lower alkyl, alkoxy, aryl, and substituted aryl.


- 20 24. The composition of claim 21 wherein R₁['], R₁^{''}, R₁^{'''}, and R₁^{''''} are each individually selected from the group H, lower alkyl, and substitute lower alkyl.

25

WO 01/40246

PCT/US00/32721

25. The composition of claim 1 wherein R^1 is selected from the group consisting of:



5 wherein each R may be independently selected from the group consisting of H, lower alkyl, and substituted lower alkyl and wherein X is O, or S (-O)₀₋₂.

26. The composition of claims 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 wherein R_1 is selected from the group consisting of 3-tetrahydrofuranyl, 3-tetrahydrothiofuranyl, 4-pyranyl, and 4 thiopyranyl.

10 27. The composition of claims 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 wherein R_1 is 3-tetrahydrofuranyl.

28. The composition of claim 1 wherein the compound is selected from the group of compounds consisting of 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-(methylthiomethyl)oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(Ethylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(Methylethylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-(phenylthiomethyl)oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(4-Methoxyphenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(4-chlorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(4-fluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(4-methylphenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(4-trifluoromethylphenylthio)methyl]oxolane-3,4-diol; 2-{6-[(3R)oxolan-3-yl]amino}purin-9-yl}{4S,5S,2R,3R)-5-[(2-Methoxyphenylthio)methyl]oxolane-3,4-diol; (5-{6-[(3R)oxolan-3-yl]amino}purin-9-yl)(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)(ethylsulfonyl)methane 2-

WO 01/40246

PCT/US00/32721

{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-[(2,4-difluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-[(2,6-dichlorophenylthio)methyl]oxolane-3,4-diol; 2-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-[(3-fluorophenylthio)methyl]oxolane-3,4-diol; 2-{6-
 5 [((3R)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-[(2-fluorophenylthio)methyl]oxolane-3,4-diol; 5-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(2S,3R,4R,5R)-4-acetyloxy-2-[(fluorophenylthio)methyl]oxolan-3-yl acetate; Methyl 2[(5-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]benzoate; {2[(5-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(2S,3S,4R,5R)-3,4-dihydroxyoxolan-2-yl)methylthio]phenyl}-N-methylcarboxamidebenzoate; 2-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-(benzoxazol-2-ylthiomethyl)oxolane-3,4-diol; 2-{6-(((3S)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-[(1-methylimidazol-2-yl-thio)methyl]oxolane-3,4-diol; 2-{6-(((3S)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-(pyrimidine-2-ylthiomethyl)oxolane-3,4-diol; 2-{6-(((3S)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-(2-pyridylthiomethyl)oxolane-3,4-diol; 2-{6-(((3S)oxolan-3-yl)amino]purin-9-yl}(4S,5S,2R,3R)-5-(4-pyridylthiomethyl)oxolane-3,4-diol; and 5-{6-(((3R)oxolan-3-yl)amino]purin-9-yl}(2S,3R,4R,5R)-4-acetyloxy-2-[(4-fluorophenylthio)methyl]oxolan-3-yl]acetate.

29. A method for modifying cardiac activity in a mammal experiencing a heart electrical disorder that can be treated by stimulating an A₁ adenosine receptor comprising the
 20 administration of a therapeutically effective amount of the composition of claim 1 to the mammal.

30. A method for modifying mammalian adipocyte function by stimulating an A₁ adenosine receptor comprising administering a therapeutically effective amount of the composition of claim 1 to the mammal.

25 31. A method to restore sensitivity and efficacy of insulin in a mammal by stimulating an A₁ adenosine receptor comprising the administration of a therapeutically effective amount of a composition of claim 1 to the mammal.

32. A method for providing a mammal with central nervous system neuroprotection by stimulating an A₁ adenosine receptor comprising administering a
 30 therapeutically effective amount of the composition of claim 1 to the mammal.

WO 01/40246

PCT/US00/32721

33. A method for providing a mammal with cardiomyocyte protection from ischemia by stimulating an A₁ adenosine receptor comprising administering a therapeutically effective amount of the composition of claim 1 to the mammal.

34. The method of claim 29 or 30 or 31 or 32 or 33 wherein the therapeutically effective amount ranges from about 0.01 to about 100 mg/kg weight of the mammal.

35. The method of claim 29 wherein the composition is administered to the mammal experiencing a heart electrical disorder selected from the group consisting of supraventricular tachycardias, atrial fibrillation, atrial flutter, and AV nodal re-entrant tachycardia.

36. The method of claim 30 or 31 wherein the composition is administered to a mammal experiencing a disorder selected from the group consisting of diabetes and obesity.

37. The method of claim 32 wherein the composition is administered to a mammal experiencing an central nervous system disorder selected from the group consisting of epilepsy, and stroke.

38. The method of claim 33 wherein the composition is administered to a mammal experiencing an ischemic event in the heart selected from the group consisting of stable angina, unstable angina, cardiac transplant, and myocardial infarction.

39. The method of claim 29 or 30 or 31 or 32 or 33 wherein the mammal is a human.

40. A pharmaceutical composition of matter comprising the composition of claim 1 and one or more pharmaceutical excipients.

41. The pharmaceutical composition of matter of claim 40 wherein the pharmaceutical composition is in the form of a solution.

42. The pharmaceutical composition of matter of claim 40 wherein the pharmaceutical composition is in the form of a tablet.

INTERNATIONAL SEARCH REPORT

National Application No PCT/US 00/32721		
A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07H19/16 A61K31/7076 A61P9/00 A61P25/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07H A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 589 467 A (LAU JESPER ET AL) 31 December 1996 (1996-12-31) claims example 21 ---	1, 2, 29, 40
A	US 5 789 416 A (LUM ROBERT T ET AL) 4 August 1998 (1998-08-04) the whole document ---	1, 29, 40
A	US 4 373 097 A (STRAMENTINOLI GIORGIO ET AL) 8 February 1983 (1983-02-08) abstract ---	1
-/--		
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents : <div style="display: flex;"> <div style="flex: 1;"> *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed </div> <div style="flex: 1;"> *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *G* document member of the same patent family </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">27 March 2001</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">05/04/2001</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">de Nooy, A</div>

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/32721

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>S. SNOWDY ET AL.: "A comparison of an A1 adenosine receptor agonist (CVT-510) with diltiazem for slowing of AV nodal conduction in guinea-pig"</p> <p>BRITISH JOURNAL OF PHARMACOLOGY, vol. 126, 1999, pages 137-146, XP000992136 abstract figure 1</p> <p>-----</p>	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

National Application No

PCT/US 00/32721

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5589467 A	31-12-1996	AU 678053 B	15-05-1997
		AU 7651994 A	03-04-1995
		CA 2171940 A	23-03-1995
		WO 9507921 A	23-03-1995
		EP 0719275 A	03-07-1996
		FI 961219 A	15-05-1996
		JP 11511436 T	05-10-1999
		NO 961071 A	15-05-1996
		NZ 273284 A	24-03-1997
		ZA 9407201 A	18-03-1996
US 5789416 A	04-08-1998	AU 726597 B	16-11-2000
		AU 4080997 A	19-03-1998
		BR 9711444 A	18-01-2000
		CN 1234803 A	10-11-1999
		CZ 9900614 A	13-10-1999
		EP 1081155 A	07-03-2001
		EP 0920438 A	09-06-1999
		EP 0992510 A	12-04-2000
		JP 2000501426 T	08-02-2000
		NO 990787 A	19-02-1999
		PL 331853 A	16-08-1999
		TR 9900377 T	21-05-1999
		WO 9808855 A	05-03-1998
US 4373097 A	08-02-1983	US 4454122 A	12-06-1984

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.